



Diet and resource partitioning between egrets and herons in an estuarine colony in southern Brazil

Amanda Oliveira Travessas · Cindy Tavares Barreto ·
Cíntia Negrine Fernandez · Daiane Carrasco Chaves ·
Leandro Bugoni

Received: 10 March 2025 / Revised: 28 January 2026 / Accepted: 30 January 2026
© The Author(s) 2026

Abstract Herons are predators at high levels of the trophic webs in aquatic and adjacent terrestrial environments. They often breed sympatrically, forming colonies with multiple species that play key roles in the transport of nutrients and can serve as environmental sentinels. However, studies evaluating foraging habits, resource partitioning, and niche

overlap are still scarce. We used complementary stable isotopes of carbon and nitrogen, and food remains analyses to investigate diet overlap among cattle egret (*Bubulcus ibis*), snowy egret (*Egretta thula*), and black-crowned night heron (*Nycticorax nycticorax*) at the Lagoa dos Patos estuary, southern Brazil. These species fed on vertebrates (mainly fish but also anurans and bird chicks) and arthropods (insects, arachnids, and crustaceans) in varying proportions. Similarly, stable isotopes demonstrated assimilated sources from estuarine, limnetic, and terrestrial areas, with food items differing in proportions. The diet of cattle egrets was composed mainly of terrestrial prey, while snowy egrets fed mainly on aquatic prey. Black-crowned night herons had a generalist diet and the highest isotopic niche breadth, which overlapped with other species. Our study demonstrated that small-to-medium-sized herons have trophic plasticity and despite sharing food items in different proportions, they show a distinction in at least one of the niche dimensions (temporal, spatial, or trophic), which allows for the coexistence of sympatric species in multispecies colonies.

Handling editor: Stuart Halse

A. O. Travessas (✉) · C. T. Barreto · C. N. Fernandez ·
L. Bugoni (✉)
Waterbirds and Sea Turtles Laboratory, Institute
of Biological Sciences, Universidade Federal do Rio
Grande - FURG, Campus Carreiros, Rio Grande,
RS 96203-900, Brazil
e-mail: amandatravessas@gmail.com

L. Bugoni
e-mail: lbugoni@yahoo.com.br

A. O. Travessas · C. N. Fernandez · L. Bugoni
Graduate Program in Biology of Continental Aquatic
Environments, Institute of Biological Sciences,
Universidade Federal do Rio Grande - FURG, Campus
Carreiros, Rio Grande, RS 96203-900, Brazil

Present Address:

C. T. Barreto
Department of Ecology and Evolutionary Biology,
University of Connecticut, Storrs, CT, USA

D. C. Chaves
Zoology Laboratory, Institute of Biological Sciences,
Universidade Federal do Rio Grande - FURG, Campus
Carreiros, Rio Grande, Rio Grande do Sul 96203-900,
Brazil

Keywords *Bubulcus ibis* · *Egretta thula* · Feeding ·
Nycticorax nycticorax · Regurgitates · Stable isotopes

Introduction

The composition of ecological communities, defined by species diversity and arrangement, is strongly influenced by abiotic and biotic factors (Begon et al., 2007; Mittelbach & McGill, 2019). The pool of n -dimensional environmental resources and conditions required by a species to exist is regarded as its niche (Hutchinson, 1957). The niche of a species can overlap among and within the niches of other species in a community (Gause, 1937). However, multiple species can still coexist under specific conditions without competing, but this coexistence depends on resource availability and strategies of the consumers (Gause, 1937; Schoener, 1974; Bearhop et al., 2004; Begon et al., 2007). Thus, studies on resource partitioning are important to understand the mechanisms that influence community structure and niche overlap (e.g., Comte et al., 2016; Faria et al., 2018; Salas-López et al., 2022; Navarro et al., 2023). Environments such as estuaries are recognized for their nutrient abundance (Day-Jr. et al., 2013) and integrate resources from marine, terrestrial, and freshwater ecosystems, providing feeding, breeding, and roosting sites for numerous bird species, including herons and egrets (Weller, 1999; Kushlan & Hancock, 2005; Kennish, 2019).

Hérons and egrets often breed in colonies with sympatric species (Kushlan & Hancock, 2005). They use environments where food resources are abundant, a strategy known as central place foraging (Bell, 1990). These birds use diverse foraging techniques related to their body and bill morphology, typically capturing their prey with their bills while wading in shallow waters (Kushlan & Hancock, 2005). Cattle egrets [*Bubulcus ibis* (Linnaeus, 1758)], snowy egrets [*Egretta thula* (Molina, 1782)], and black-crowned night herons [*Nycticorax nycticorax* (Linnaeus, 1758)] are sympatric, small-to-medium-sized, predatory waterbirds (Kushlan & Hancock, 2005; Hothem et al., 2020; Parsons & Master, 2020; Telfair-II, 2023) and, therefore, with potential niche overlap and/or niche segregation.

The cattle egret is originally from Africa (Ducommun et al., 2008; Moralez-Silva & Del Lama, 2014) but now has colonized the New World due to its great capacity to invade and occupy new areas (Kushlan & Hancock, 2005; Nunes et al., 2010; Abdullahm et al., 2017). The species

inhabits grasslands and benefits from insects that emerge from the soil with cattle movement (Bó & Darrieu, 1993; Kaufman, 1996; Kushlan & Hancock, 2005). Its variable diet includes insects, arachnids, and amphibians (Sánchez-García, 2012; Vega-Sánchez et al., 2022). The snowy egret has a wide distribution, occurring in aquatic environments throughout the Americas (Martínez-Vilalta & Motis, 1992; Brzorad & Maccarone, 2013). The species feeds mainly on fish, but crustaceans, amphibians, reptiles, and insects are also among their prey (Smith, 1997; Antón-Tello et al., 2021). The black-crowned night heron occurs in the Americas, Europe, Africa, and Asia (Hancock & Kushlan, 2010). It is considered an opportunistic species, foraging solitarily during the night or twilight (Kushlan & Hancock, 2005; Maccarone et al., 2015). As observed for cattle and snowy egrets, black-crowned night herons also have a varied diet, consuming items such as fish, amphibians, reptiles, small rodents, bats, insects, spiders, crustaceans, mollusks, birds (including cannibalism), and eggs of many other species (Kazantzidis & Goutner, 2005; Kushlan & Hancock, 2005; Riehl, 2006; Régale et al., 2014). These three species are often seen using habitats modified by humans such as artificially created vegetated banks, shallow freshwater environments used for irrigation of crops, saltwater rivers, man-made streams, lakes, and lagoons, and anthropically modified swamps and mangroves (Kushlan & Hancock, 2005).

Despite their substantial ecological influence in estuaries, where they facilitate nutrient transfer from aquatic to terrestrial ecosystems, subsidizing trophic interactions (Caseiro-Silva et al., 2023), detailed information on the diet and ecological roles of herons and egrets in these environments remains scarce. To understand the role of cattle egrets, snowy egrets, and black-crowned night herons in estuaries, their interspecific interactions and resource partitioning, food preferences, and energy and nutrient flow should be elucidated (Steinmetz et al., 2003; Green & Elmberg, 2014). These can be assessed through diet and isotopic niche studies.

Conventional diet studies can be applied and allow us to determine and quantify the prey consumed through the fragments found in regurgitates and pellets. Regurgitates represent partially

digested prey, while pellets are composed of hard remnants of undigested food items covered by mucus. Both regurgitates and pellets allow for estimates of the contribution of different food items in a consumer's diet, associated with niche metrics. These analyses provide essential background information for more advanced research techniques, such as stable isotopic analyses, enhancing the accuracy of isotopic mixture models (Layman et al., 2012; Parnell et al., 2013; Britto & Bugoni, 2015).

Carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) stable isotopes are commonly used to identify food resources assimilated in consumer tissues. This technique provides valuable insights into the prey species consumed, isotopic niche overlap, and isotopic niche breadth, allowing for a comparison across various trophic niche dimensions (Newsome et al., 2007; Layman et al., 2012; Hette-Tronquart, 2019). While $\delta^{13}\text{C}$ is particularly useful in identifying environmental sources of food, as it considers the photosynthetic cycle of primary C3 vs C4 producers, leading to a slight variation in consumer-tissue discrimination, $\delta^{15}\text{N}$ is more suitable for identifying trophic relationships, as it increases at each trophic level (Post, 2002; Staddon, 2004; Fry, 2006; Castro et al., 2016).

In this study, we examined trophic niche partitioning among three small-to-medium-sized estuarine herons and egrets that breed sympatrically, through complementary techniques, i.e., analyses of regurgitates, pellets, and stable isotopes. We hypothesized that cattle egrets would have a distinct niche as they forage mainly in terrestrial environments, compared to the other two species that feed mainly in aquatic environments. We further hypothesized that snowy egrets would have a restricted niche as they forage preferentially in coastal, estuarine, and limnetic environments. Finally, we hypothesized that black-crowned night herons would have the most diverse diet and high trophic and isotopic niche overlap. As cattle egrets have a predominantly terrestrial feeding habit, foraging primarily on insects available nearby nests, we expect that cattle egrets will occupy a lower trophic level in comparison to snowy egrets and black-crowned night herons, which are species that feed predominantly on carnivorous fish and crustaceans available in the aquatic environments. Conversely, we expect to find a higher trophic level and wider isotopic niche for black-crowned night herons


compared to the egrets, reflecting their opportunistic feeding habits, which include animals of high trophic levels, such as predatory fish, birds, amphibians, and mammals.

Methods

Study area and field sampling

Bird sampling took place on Marinheiros Island (32°00' S, 52°09' W; Fig. 1), with 40 km², in the Lagoa dos Patos Estuary, Rio Grande city, Rio Grande do Sul, Brazil. The island is surrounded by a belt of wetland environments and composed of horticultural areas, swamp ('restinga') forests in the peripheral zone, dunes, and a lagoon in the center of the island (Gianuca et al., 2010). The climate in the region is temperate (Klein, 1997), with a variation of ambient temperatures from 5.9 to 35 °C during the spring of 2019 and from 12.8 to 35.9 °C in the summer of 2020 (INMET, 2023). The salinity and water level are influenced by climate and rainfall, with the highest salinity consistently occurring in summer (Garcia et al., 2001). The island is situated in an important region for the conservation of waterbirds (Williamson et al., 2013), and a colony of Pelecaniformes is established yearly in the restinga forest, a few hundred meters from the shore of Lagoa dos Patos (Gianuca et al., 2010; Britto & Bugoni, 2015; Barreto et al., 2025). This colony is the breeding site of two species of Threskiornithidae, the roseate spoonbill [*Platalea ajaja* Linnaeus, 1758 and the bare-faced ibis, *Phimosus infuscatus* (Lichtenstein, 1823)], and seven species of Ardeidae, the great egret (*Ardea alba* (Linnaeus, 1758)], the cocoi heron (*Ardea cocoi* (Linnaeus, 1766)], the cattle egret, the little blue egret [(*Egretta caerulea* (Linnaeus, 1758)], the snowy egret, the yellow-crowned night heron [(*Nyctanassa violacea* (Linnaeus, 1758)], and the black-crowned night heron (Gianuca et al., 2010; Barreto et al., 2025). The trees in which each species will nest in a particular year, as well as the nest height, seems to be related to arrival of breeding pairs, but does not have a particular species-specific pattern in multispecies colonies, although there is a tendency of bigger species (i.e., spoonbills, cocoi herons, and great egrets) arriving earlier and, therefore, nesting in

Fig. 1 Marinheiros Island, Lagoa dos Patos Estuary, southern Brazil. The star indicates the colony location where cattle egret (*Bubulcus ibis*), black-crowned night heron (*Nycticorax nycticorax*), and snowy egret (*Egretta thula*) breed annually. (Map courtesy of A.B. da Silva)



taller trees (Bachir et al., 2008, Gianuca et al., 2010; Park et al., 2011).

Spontaneous regurgitates and pellets were sampled from chicks of three ardeids (23 black-crowned night herons, six cattle egrets, and three snowy egrets) inside the nests or their surroundings, during the breeding season, between September 2019 and January 2020, for conventional analysis of the diet. Dietary samples were stored in plastic bags and frozen until analysis. For stable isotope analysis sampling, chicks were manually captured, morphometric measurements were taken to infer age, and approximately 0.1 mL of blood was drawn from their tarsal vein by using a syringe and needle. To avoid resampling, birds were banded with uniquely numbered metal bands provided by the Centro Nacional de Pesquisa e Conservação de Aves Silvestres (CEMAVE) of the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio). Blood samples were preserved in 1.5 mL vials in the field, taken to the Waterbirds and Sea Turtles Laboratory (LAATM) of the

Universidade Federal do Rio Grande (FURG), and frozen until analysis. Only chicks at least 3 weeks old were sampled, as it is expected that due to the turnover time range of 7–20 days (Boecklen et al., 2011), the isotopic composition of the blood will represent values of the food delivered by parents rather than egg nutrients. A total of 43 samples from different individuals were collected for stable isotope analysis (15 cattle egrets, 20 black-crowned night herons, and eight snowy egrets). Additionally, insects of the orders Orthoptera, Odonata, and Coleoptera and spiders, to represent the isotopic values of consumed invertebrates, were collected manually. Finally, whole or minimally digested fish present in the regurgitates had tissues sampled to represent the isotopic values of limnetic and estuarine fish consumed.

Conventional diet data analysis

To identify the consumed prey, items found in the samples were separated by washing the regurgitates

and pellets under running water in a 0.75 mm mesh metal sieve. Intact prey items were separated from other food items (parts of prey items). All consumed food items were quantified (i.e., counted), measured in millimeters with a caliper, and weighed in grams on a laboratory scale. For identification, items were examined under a stereomicroscope, and prey items were compared with reference collections and identification guides (Fisher et al., 2004; Malabarba et al., 2013; Conversani et al., 2017; Hamada et al., 2018).

For each prey species, the relative frequency of occurrence (FO%), which is the presence of the taxon related to the total number of samples, the mean percentage contribution of each food item (PN%), and the percentage contribution of the reconstituted mass of each food item (PM%) were calculated. Prey mass was reconstructed using allometric equations based on measurements of fish otoliths or from mean values of similar-sized prey (Bugoni & Vooren, 2004). Due to the low sample size of snowy egrets, dietary parameters were calculated only for cattle egrets and black-crowned night herons. After obtaining FO%, PN%, and PM%, we calculated the prey-specific relative importance index (PSIRI%) according to Brown et al. (2012):

$$\text{PSIRI}\% = [\text{FO}\% \times (\text{PN}_i\% + \text{PM}_i\%)]/2 \quad (1)$$

where FO% is the frequency of occurrence of each taxon; PN_i% is the contribution in number of each taxon; and PM_i% is the mass contribution of each taxon.

To calculate the trophic niche overlap among species, based on the contribution in numbers of each taxon (N%), we used the Morisita-Horn similarity index (Horn, 1966), following the equation:

$$\text{CH} = [2(\sum ni \times p_{ij} \times p_{ik})]/(\sum ni \times p_{2ij} + \sum ni \times p_{2ik}) \quad (2)$$

where CH is the Morisita-Horn similarity index; *n* is the total number of resources used; *p_{ij}* is the proportion of resources *i* to the total resources used by species *j* and *p_{ik}* is the proportion of resource *i* on the total resources used by species *k*.

For the trophic niche breadth, which is based on the contribution in number of each taxon, we used the Levin's standardized index (Magurran, 2019), following the equation of Hurlbert (1978):

$$B_s = [(\sum p_i^2) - 1]/(n-1) \quad (3)$$

where *B_s* is Levin's standardized index; *p_i²* is the proportion of the resource belonging to resource category *i*; and *n* is the total number of food items consumed.

Stable isotope analysis (SIA)

In the laboratory, whole blood samples were freeze-dried, homogenized, and weighed in tin capsules. Insects, spiders, crabs, anurans, and fish samples from the regurgitates were placed in a Soxhlet extractor with petroleum ether for 6 h for lipid removal (Post et al., 2007; Elliott et al., 2017; Cloyd et al., 2020). Subsequently, the prey samples were freeze-dried, homogenized, and weighed in tin capsules. Blood and prey samples were then analyzed with an isotope ratio mass spectrometer (IRMS) coupled to an elemental mass analyzer at the Integrated Analysis Center at FURG (CIA-FURG, Brazil). The isotopic signature values are expressed by the delta notation (δ), in parts per thousand (‰), and differences between the stable isotope values of the sample compared to standards are given by Eq. 4, from Bond & Hobson (2012):

$$\delta^{13}\text{C} \text{ or } \delta^{15}\text{N} (\text{‰}) = (R_{\text{sample}}/R_{\text{standard}}) - 1 \quad (4)$$

where δ¹³C is the carbon stable isotope; δ¹⁵N is the nitrogen stable isotope; *R_{sample}* is the ratio between the heavy and light isotopes of the sample; and *R_{standard}* is the ratio between the heavy and light isotopes of the standard. Glutamic acid, caffeine, and acetazolamide were the internal standards at CIA, and Vienna Pee Dee Belemnite limestone and air were carbon and nitrogen, respectively, as calibration standards. The internal standards had a deviation of 0.1‰ for δ¹³C and 0.4‰ for δ¹⁵N. For prey taxonomic groups not found in regurgitated samples nor sampled at the colony, isotopic data from the LAATM-FURG isotopic database were used.

Bayesian isotopic mixing models were generated on 'simmr' R package (Parnell et al., 2013) in R software version 4.0.1 (R Core Team, 2020) to estimate the contribution of food items in the whole blood of chicks. For the models, the most common prey found in regurgitates and pellets were used. The trophic discrimination factors (TDF) of the consumers in the models were $-0.3 \pm 0.5\text{‰}$ and $2.61 \pm 0.5\text{‰}$

for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, respectively, as they were used in previous studies with isotopic modeling at the same or closely related waterbirds (Silva-Costa & Bugoni, 2013; Britto & Bugoni, 2015; Faria et al., 2016). These TDFs were obtained from experimental studies with penguins, which are predominantly piscivorous birds that feed on whole fish [king penguin *Aptenodytes patagonicus* J.F. Miller, 1778: $\delta^{13}\text{C} = -0.81\text{‰}$ and $\delta^{15}\text{N} = 2.07\text{‰}$; southern rockhopper penguin *Eudyptes chrysocome* (J.R. Forster, 1781): $\delta^{13}\text{C} = 0.20\text{‰}$ and $\delta^{15}\text{N} = 2.72\text{‰}$; Chérel et al., (2005)], and from experiments with tufted puffins [*Fratercula cirrhata* (Pallas, 1769): $\delta^{13}\text{C} = -0.30$ and $\delta^{15}\text{N} = 3.05\text{‰}$; Williams et al., (2007)]. Finally, standard ellipse areas adjusted for small sample sizes (SEAc) were used to determine the isotopic niche breadth and overlap of isotopic niches of the three consumers using the stable isotope Bayesian ellipses in R (Jackson et al., 2011).

Results

Conventional diet analysis and trophic niche characterization

The analysis of regurgitates and pellets indicates a cattle egret diet composed mainly of insects, spiders, and anurans, but fish, crustaceans, and ticks (Table 1). The most frequent food items were insects (FO% = 100), which had an importance (PSIRI%) of 43.2, followed by anurans (FO% = 83.3%; PSIRI% = 26.6), and arachnids (FO% = 83.3%; PSIRI% = 13.0). Anurans and insects had the highest contribution, according to the percentage of their reconstituted mass (PM%), in the diet of cattle egrets (52.1% and 27.4%, respectively). The trophic niche breadth was $B_s = 0.12$.

The conventional analysis indicates a variety of items composing the diet of black-crowned night herons, which consists of insects, crustaceans, fish, spiders, tadpoles, and heron chicks, including cannibalism (Table 1). Insects (FO% = 78.2) and fish (FO% = 73.9) were the most frequently occurring prey. Notwithstanding, the highest contributions to reconstituted mass were bird chicks (PM% = 44.1), followed by fish (PM% = 29.1), while the most

Table 1 Mean values of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ (‰ ± standard deviation) in the blood of chicks of the cattle egret (*Bubulcus ibis*), the black-crowned night heron (*Nycticorax nycticorax*), and the

snowy egret (*Egretta thula*), and muscle/whole individual of potential prey from southern Brazil, used in the Bayesian mixing models. n = the number of samples analyzed

Birds and potential sources	Species/taxon	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	n	References
Chick blood	<i>Bubulcus ibis</i>	-22.47 ± 2.53	7.17 ± 0.78	15	This study
	<i>Egretta thula</i>	-23.77 ± 2.26	8.25 ± 0.35	8	This study
	<i>Nycticorax nycticorax</i>	-22.87 ± 2.64	9.91 ± 1.91	20	This study
Anurans		-23.31 ± 1.06	4.22 ± 0.82	5	This study
	<i>Pseudis minuta</i>	-24.77	5.18	1	This study
	<i>Scinax squaleirostris</i>	-23.90	4.32	1	This study
	<i>Physalaemus gracilis</i>	-22.63 ± 0.71	3.87 ± 0.83	3	This study
Estuarine fishes	<i>Atherinella brasiliensis</i>	-14.61	12.29	2	Garcia et al. (2007)
Freshwater fishes	<i>Astyanax</i> sp.	-20.25 ± 1.9	11.82 ± 0.3	2	Garcia et al. (2007)
Crustaceans	<i>Callinectes</i> sp.	-18.68 ± 0.14	8.36 ± 1.63	2	This study
Aquatic insects	Belostomatidae	-28.3	5.7	1	Britto & Bugoni (2015)
Terrestrial insects		-25.67 ± 1.91	5.89 ± 1.96	6	
	Coleoptera	-25.65 ± 2.97	5.97 ± 2.20	3	This study
	Odonata	-25.34	5.09	1	LAATM database ^a
	Orthoptera	-25.88 ± 0.66	6.17 ± 2.96	2	This study and LAATM database

^a Unpublished data

Table 2 Diet composition of chicks of the cattle egret (*Bubulcus ibis*) and the black-crowned night heron (*Nycticorax nycticorax*) at the colony of Marinheiros Island, Lagoa dos Patos Estuary, Brazil

Food items	Cattle egret (n=6)				Black-crowned night heron (n=23)			
	FO%	PN%	PM%	PSIRI%	FO%	PN%	PM%	PSIRI%
Anurans	83.3	12.4	52.1	26.6	4.3	5.8	0.06	0.1
Hylidae	66.6	9.7	41.7	17.0	–	–	–	–
<i>Hypsiboas pulchellus</i>	16.6	0.3	0.3	<0.1	–	–	–	–
<i>Pseudis minuta</i>	33.3	5.8	14.3	3.3	–	–	–	–
<i>Scinax squalirostris</i>	33.3	3.7	27.1	5.0	–	–	–	–
Leiuperidae	50.0	2.1	1.6	0.9	–	–	–	–
<i>Physalaemus gracilis</i>	33.3	1.8	1.4	0.5	–	–	–	–
<i>Pseudopaludicola falcipes</i>	16.6	0.3	0.2	<0.1	–	–	–	–
Leptodactylidae	33.3	0.5	8.8	1.5	–	–	–	–
<i>Leptodactylus gracilis</i>	16.6	0.3	0.7	<0.1	–	–	–	–
<i>Leptodactylus latrans</i>	16.6	0.3	8.0	0.7	–	–	–	–
Unidentified	–	–	–	–	4.3	5.8	0.06	0.1
Fishes	16.6	5.3	4.8	0.8	73.9	12.1	29.1	15.2
Atherinopsidae	–	–	–	–	30.4	3.9	23.1	4.1
<i>Atherinella brasiliensis</i>	–	–	–	–	30.4	3.9	23.1	4.1
Characidae	–	–	–	–	8.7	0.8	0.2	<0.1
<i>Astyanax</i> sp.	–	–	–	–	4.3	0.4	0.2	<0.1
<i>Pseudocorynopoma doriae</i>	–	–	–	–	4.3	0.4	0.04	<0.1
Poeciliidae	16.6	5.3	4.8	0.8	8.7	0.8	0.1	<0.1
<i>Phalloceros caudimaculatus</i>	16.6	5.3	4.8	0.8	8.7	0.8	0.1	<0.1
Cichlidae	–	–	–	–	4.3	0.4	0.09	<0.1
Sciaenidae	–	–	–	–	4.3	0.4	0.3	<0.1
<i>Stellifer</i> sp.	–	–	–	–	4.3	0.4	0.3	<0.1
Ariidae	–	–	–	–	4.3	0.8	0.8	<0.1
<i>Genidens barbatus</i>	–	–	–	–	4.3	0.8	0.8	<0.1
Loricariidae	–	–	–	–	8.7	0.8	0.8	<0.1
<i>Loricariichthys anus</i>	–	–	–	–	8.7	0.8	0.8	<0.1
Synbranchidae	–	–	–	–	8.7	0.8	1.0	<0.1
<i>Synbranchus marmoratus</i>	–	–	–	–	8.7	0.8	1.0	<0.1
Unidentified	–	–	–	–	17.4	2.3	2.8	0.4
Birds	–	–	–	–	8.7	0.8	44.1	1.9
Ardeidae	–	–	–	–	8.7	0.8	44.1	1.9
<i>Bubulcus ibis</i> (chicks)	–	–	–	–	4.3	0.4	20.7	0.4
<i>Nycticorax nycticorax</i> (chicks)	–	–	–	–	4.3	0.4	23.4	0.5
Insects	100.0	58.9	27.4	43.2	78.3	57.0	5.2	24.3
Blattaria	50.0	2.9	1.4	2.6	4.3	0.4	0.04	<0.1
Blattidae	16.6	1.3	0.2	0.1	–	–	–	–
Coleoptera	66.6	13.9	2.1	5.3	56.5	41.0	3.8	12.6
Coccinellidae	16.6	0.3	0.2	<0.1	–	–	–	–
<i>Eriopsis connexa</i>	16.6	0.3	0.2	<0.1	–	–	–	–
Hydrophilidae	–	–	–	–	4.3	0.4	0.04	<0.1
Scarabaeidae	–	–	–	–	4.3	0.4	0.04	<0.1
Diptera	66.6	15.3	9.3	8.1	–	–	–	–
Brachycera	66.6	7.9	5.1	4.3	–	–	–	–

Table 2 (continued)

Food items	Cattle egret (<i>n</i> =6)				Black-crowned night heron (<i>n</i> =23)			
	FO%	PN%	PM%	PSIRI%	FO%	PN%	PM%	PSIRI%
Dolichopodidae	16.6	0.5	0.3	<0.1	–	–	–	–
Muscidae	16.6	0.5	0.3	<0.1	–	–	–	–
Sarcophagidae	33.3	5.0	3.0	1.3	–	–	–	–
Tabanidae	16.6	0.8	0.5	0.1	–	–	–	–
Nematocera	16.6	0.5	0.3	<0.1	–	–	–	–
Tipulidae	16.6	0.5	0.3	<0.1	–	–	–	–
Hemiptera	50.0	2.6	1.4	1.0	39.1	6.2	0.6	2.6
Auchenorrhyncha	–	–	–	–	21.7	2.7	0.2	0.3
Cercopidae	–	–	–	–	8.7	1.6	0.1	<0.1
Heteroptera	33.3	1.0	0.6	0.3	17.3	1.9	0.2	0.18
Belostomatidae	33.3	0.5	0.3	0.1	13.0	1.6	0.1	0.1
<i>Belostoma</i> sp.	16.6	0.3	0.2	<0.1	4.3	0.8	0.07	<0.1
Pentatomidae	16.6	0.5	0.3	<0.1	–	–	–	–
Naucoridae	–	–	–	–	4.3	0.4	0.04	<0.1
Hymenoptera	50.0	1.0	0.8	0.5	30.4	7.0	0.6	1.1
Formicidae	16.6	0.3	0.2	<0.1	8.7	0.8	0.07	<0.1
<i>Camponotus</i> sp.	–	–	–	–	4.3	0.4	0.04	<0.1
<i>Solenopsis invicta</i>	16.6	0.3	0.3	<0.1	–	–	–	–
Neuroptera	33.3	0.8	0.6	0.2	–	–	–	–
Odonata	33.3	2.4	1.4	0.6	4.3	1.6	0.1	<0.1
Coenagrionidae	16.6	0.5	0.3	<0.1	–	–	–	–
Libellulidae	33.3	1.8	1.3	0.5	4.3	1.6	0.1	<0.1
Orthoptera	66.6	20.0	12.2	10.6	4.3	1.2	0.1	<0.1
Acrididae	50.0	3.9	2.4	1.6	–	–	–	–
<i>Aleuas</i> sp.	16.6	0.3	0.2	<0.1	–	–	–	–
Gryllidae	50.0	7.4	4.5	3.0	–	–	–	–
Tettigoniidae	33.3	5.3	1.6	1.1	–	–	–	–
Romaleidae	16.6	0.5	0.3	<0.1	4.3	1.2	0.1	<0.1
<i>Alcamenes clarazianus</i>	16.6	0.5	0.3	<0.1	4.3	1.2	0.1	<0.1
Arachnids	83.3	16.6	14.7	13.0	17.3	6.6	0.9	0.6
Araneae	83.3	15.5	14.3	12.4	17.3	6.6	0.9	0.6
Ixodidae	33.3	1.0	0.4	0.2	–	–	–	–
<i>Rhipicephalus sanguineus</i>	16.6	0.8	0.3	0.1	–	–	–	–
Crustaceans	33.3	1.6	1.0	0.4	21.7	17.6	20.6	4.1
Decapoda	–	–	–	–	13.0	4.3	18.1	1.4
Grapsidae	–	–	–	–	4.3	0.4	0.4	<0.1
Portunidae	–	–	–	–	8.7	1.6	17.8	0.8
<i>Callinectes</i> sp.	–	–	–	–	8.7	1.6	17.8	0.8
Isopoda	33.3	1.6	1.0	0.4	8.7	13.3	1.2	0.6
Ligiidae	33.3	1.6	1.0	0.4	8.7	13.3	1.2	0.6
<i>Ligia exotica</i>	33.3	1.6	1.0	0.4	8.7	13.3	1.2	0.6
		<i>n</i> =380				<i>n</i> =256		

FO%=frequency of occurrence; PN%=contribution in prey-specific numbers; PM%=contribution in prey-specific reconstituted mass; PSIRI%=prey-specific relative importance index. *n*=the number of samples analyzed

important prey were insects (PSIRI% = 24.3) and fish (PSIRI% = 15.2). The trophic niche breadth was $B_s = 0.63$. The niche overlap between black-crowned night herons and cattle egrets was high ($CH = 0.67$; Table 2), demonstrating an elevated overlap.

Snowy egret ($n = 3$) regurgitates indicated a diet composed of two killifish (i.e., small freshwater annual fish) species [*Phalloceros caudimaculatus* (Hensel, 1868) and *Cnesterodon decemmaculatus* (Jenyns, 1842)], three unidentified fish, and two insect groups (Belostomatidae and Odonata). Due to the low sample size, snowy egrets were not considered in the niche overlap analysis.

Prey assimilation and isotopic niche

Values for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in the blood of black-crowned night herons were more variable in comparison to the other two species, while values from snowy egret chicks were less variable (Table 1; Fig. 2). Mean $\delta^{13}\text{C}$ value in the blood of snowy egrets was -23.5‰ , while the mean value for cattle egrets was -22.7‰ and for black-crowned night herons -22.9‰ . The highest values of $\delta^{15}\text{N}$, on the other hand, were from the blood of black-crowned night heron chicks (9.9‰), followed by blood from snowy egret chicks (8.2‰), and cattle egrets (7.2‰).




Fig. 2 Individual values of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ (‰) in chick blood of cattle egret (*Bubulcus ibis*), black-crowned night heron (*Nycticorax nycticorax*), and snowy egret (*Egretta thula*). The means \pm SD of isotopic values in their prey are from the estuarine environment, represented by estuarine fish (i.e., the Brazilian silverside *Atherinella brasiliensis*) and crustaceans

As observed in the reconstituted mass analysis of cattle egrets ($n = 15$), stable isotopes mixing models indicated that anurans were their main food source (95% CI—credible interval = 10–90), followed by terrestrial insects (95% CI 0.8–21; Fig. 3a). For black-crowned night herons ($n = 20$), the stable isotopes mixing models indicated anurans (95% CI 4–57) and terrestrial insects (95% CI 2.4–55) as the resources most assimilated (Fig. 3b), contrasting the reconstituted mass results (from the conventional analysis of regurgitates) for the species. Finally, for snowy egrets ($n = 8$), anurans (95% CI 11–73) and aquatic insects (95% CI 2.8–60) were the most assimilated prey sources (Fig. 3c).

Regarding the isotopic niche (Table 2; Fig. 4), black-crowned night herons had the largest niche breadth ($SEAc = 93.4$), followed by cattle egrets ($SEAc = 29.6$), and snowy egrets with the lowest niche breadth ($SEAc = 7.8$). The isotopic niche of cattle egrets overlapped substantially (73.8%) with black-crowned night herons, but only 17% with snowy egrets. The niche of snowy egrets, on the other hand, overlapped only 8.3% with black-crowned night herons. The most surprising result was for the niche area of black-crowned night herons, which overlapped 100% with the niche area of snowy egrets and 73.8% with cattle egrets.

(*Callinectes* sp.). Prey from the limnetic environment are represented by freshwater fishes (*Astyanax* sp.) and aquatic insects (Belostomatidae). Terrestrial prey are represented by anurans (i.e., *Pseudis minuta*, *Physalaemus gracilis*, and *Scinax squarilirostris*) and terrestrial insects (Coleoptera, Odonata, and Orthoptera)

Fig. 3 Contribution of potential prey sources (AN = anurans, EF = estuarine fishes, FF = freshwater fishes, AI = aquatic insects, TI = terrestrial insects, and CR = crustaceans) as evidenced in the chick blood of (a) cattle egret (*Bubulcus ibis*), (b) black-crowned night heron (*Nycticorax nycticorax*), and (c) snowy egrets (*Egretta thula*) in an estuarine colony of Pelecaniformes on Marinheiros Island, Lagoa dos Patos Estuary, southern Brazil. The horizontal line in the center of the rectangle represents the mean, and the rectangle indicates the standard deviation. The vertical lines outside the rectangle indicate the minimum and maximum contribution values





Fig. 4 Isotopic niche of cattle egret (*Bubulcus ibis*), black-crowned night heron (*Nycticorax nycticorax*) and snowy egret (*Egretta thula*) in delta space (δ) on Marinheiros Island, Lagoa dos Patos Estuary, southern Brazil, obtained through corrected standard ellipse areas (SEAc). The inset shows SEAc at $\%o^2$ based on isotopic values in whole blood



Discussion

The three ardeids studied in southern Brazil had substantial overlap in their diet and trophic niches. Anurans were the most assimilated sources for all three species according to SIA, while insects were the most important food items in the diet of cattle egrets and black-crowned night herons according to pellet and regurgitation analysis. As demonstrated by $\delta^{15}\text{N}$ values, cattle egrets were at the lowest trophic level, followed by snowy egrets, and black-crowned night herons, matching our predictions. The diet of cattle egrets was mainly composed by terrestrial insects, matching our hypothesis of a terrestrial niche for the species. Snowy egrets had a more aquatic diet, composed mainly of fish and aquatic insects, which matches our hypothesis of a more restricted aquatic niche for the species. Black-crowned night herons had the most generalist feeding habit and most diverse diet, as well as the largest overlap of trophic niche with cattle egrets and isotopic niche with cattle and snowy egrets, marching our hypothesis of high trophic level and extensive niche overlap.

The Lagoa dos Patos Estuary is a relevant area for the conservation of Pelecaniformes in South America (Kushlan & Hancock, 2005; Dias et al., 2017) and has a high abundance of species that serve as food resources for birds (Pereira & D’Incao, 2012; Haimovici & Cardoso, 2016; Teixeira-Amaral et al., 2017; Lemos et al., 2022). The conventional diet analysis provided detailed information about the prey consumed by small herons and egrets from Marinheiros

Island, similar to findings of other Pelecaniformes species in the same colony (Britto & Bugoni, 2015). Pellets and regurgitates, therefore, proved to be a reliable technique to determine the differences in food resources used by sympatric waterbird species breeding in heterogeneous environments.

The mainly terrestrial diet of cattle egrets diet corroborates with other studies (Ashoori et al., 2017a, b; Antón-Tello et al., 2021), predominantly insects, but also with varied taxa (Diptera, Orthoptera, and Hemiptera), in addition to small anurans and spiders, indicating that terrestrial environments near the colony were the main foraging areas for cattle egrets. Aquatic prey, including killifishes, and semiaquatic prey, such as giant waterbugs, were also present in the cattle egret diet, although consumed less frequently. Our results point to an opportunistic feeding habit, demonstrating that cattle egrets can be highly adaptive and explore both aquatic and terrestrial environments, similarly to the opportunistic feeding habits previously found (Kioko et al., 2016). Regarding the assimilation of consumed food items, isotopic results differed from a study in Spain (Antón-Tello et al., 2021), which showed insects as the main assimilated prey items for cattle egrets, followed by anurans. At Lagoa dos Patos, cattle egrets had anurans as the predominant assimilated food source, followed by terrestrial insects. Our models also indicated that cattle egrets have narrow trophic and isotopic niches due to the exploitation of predominantly terrestrial prey. This supports our hypothesis that the species has a

distinct niche because it forages primarily in terrestrial environments associated with cattle.

Black-crowned night herons also have a generalist feeding habit, matching our hypothesis of foraging in terrestrial, limnetic, and estuarine environments, with insects (Coleoptera, Hemiptera, and Hymenoptera) and fish of different sizes, such as the Brazilian silverside [*Atherinella brasiliensis* (Quoy & Gaimard, 1825)], as their main prey. Insects are among the most consumed prey items by black-crowned night herons elsewhere (e.g., Kazantzidis & Goutner, 2005), but fish are the most important prey in wetland areas (e.g., Ashoori et al., 2017a, b; Dias et al., 2025) as amphibians are in rice plantations (e.g., Cardarelli et al., 2017). Fish is an important prey for other Pelecaniformes breeding nearby the Lagoa dos Patos Estuary (Britto & Bugoni, 2015; Faria et al., 2016), and a variety of species of different sizes are present in both estuarine and limnetic areas in or near our study site (Quintela et al., 2018). Heron chicks, tadpoles, crustaceans, and spiders were also among the important food items consumed by black-crowned night herons in our study. Cannibalism by adults and juveniles feeding on chicks, which we detected in our samples, is reported in other studies (e.g., Riehl, 2006; Brussee et al., 2017; Sovrano et al., 2022), confirming the opportunistic behavior of the species (Davis-Jr., 1993). Regarding the taxa we found in the diet of black-crowned night herons, they were previously reported in South America in the Paraná River, Argentina (Quiroga et al., 2013). Our study found a more diverse diet of black-crowned night herons compared to other studies worldwide (review in Hothem et al., 2020), which can be explained by their opportunistic feeding behavior (Kazantzidis & Goutner, 2005) and foraging habits in varied environments (Dias et al., 2025). The SIA demonstrated that black-crowned night herons assimilated more sources from estuarine environments, such as fish and crustaceans, in addition to other sources from terrestrial and limnetic environments, distinct from the much narrower isotopic niche reported in a mixed colony at the New York/New Jersey Harbor Estuary (Craig et al., 2015). Results presented here confirmed our hypothesis that black-crowned night herons had the widest range of trophic and isotopic niches compared to egrets, as well as the greatest variety of taxa preyed upon.

For snowy egrets, two freshwater killifishes, plus Odonata and Belostomatidae, both aquatic insects,

were the main consumed food items, which matches our hypothesis that the species would have a more aquatic diet and niche. Even with limited sampling, the results were similar to those of previous studies in coastal limnetic environments (Martínez, 2010; Ruiz-Guerra & Echeverry-Galvis, 2019). Another study at the same area demonstrated similar results, as snowy egrets tended to use limnetic environments more than other heron and egret species (Gianuca et al., 2010). Fishes and crustaceans are also important prey in a previous study in the area (Gianuca et al., 2010) as well as in the upper Paraná River floodplain (Dias et al., 2025).

Snowy egrets in our study had greater assimilation of limnetic fish, showing little variation in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in blood. These results match our hypothesis of a more restrictive niche, which demonstrates that snowy egrets forage in areas near colonies rather than on marine beaches a few kilometers away. They are capital breeders, as shown by stable isotope values in eggs in a Colorado River colony in Mexico (Herzka et al., 2013). The specialist habitat of the snowy egret led to a restrict use of resources, when compared to the other species in our study. Regarding the overlap of the trophic niches between cattle egrets and black-crowned night herons, the consumption of common prey in terrestrial environments, such as insects and spiders, may characterize the most shared resources between the species. Even with the small sample size of the snowy egret, the presence of the insects in its diet was noted. This fact demonstrates that insects are important food items for all three heron species. Therefore, the current decline in the abundance and diversity of insects (Wagner et al., 2021) and the decrease in the abundance of the Brazilian silverside in the Lagoa dos Patos Estuary (Silva et al., 2021) point toward a reduction in key prey for egrets and herons, potentially influencing the whole island ecosystem. The black-crowned night heron also had a high isotopic niche overlap with the snowy egret (100%), followed by overlap with the cattle egret's niche (73.8%), which can be explained by the assimilation of varied sources, such as fish species, in comparison to other ardeids. This generalist feeding habit had led to a broader niche and resulting in high overlap with the other species. Herons and egrets that breed in sympatry may show a distinction in at least one of the niche dimensions (temporal, spatial, or trophic) during the reproductive period, which allows

for coexistence (Ye et al., 2021), as confirmed in our study.

Conclusion

Results showed that cattle egrets, snowy egrets, and black-crowned night herons share a range of upper-level prey taxa, yet differed in prey species assimilated. Although these species forage in distinct habitats from one another, they share prey species, resulting in trophic and isotopic niche overlaps. Therefore, the differences in prey consumed and assimilated and the exploration of distinct foraging sites at different periods of the day seem to allow the sympatric coexistence of the species.

The present study is the first to compare, with conventional and SIA methods, the diet and investigate resource partitioning among three ardeid species (*Bubulcus ibis*, *Egretta thula* and *Nycticorax nycticorax*) breeding at Lagoa dos Patos area. Consequently, it is important to obtain more in-depth information, especially about trophic ecology, as we know that herons, egrets, and ibises are important aquatic-terrestrial carriers of matter and nutrients (Caseiro-Silva et al., 2023) and environmental sentinels (Barreto et al., 2025). Additionally, Lagoa dos Patos is an important site for a range of waterbird species (Gianuca et al., 2010; Dias et al., 2017), emphasizing the importance of collecting in situ ecological information. Further understanding of the consumer-prey relationships, movements between environments, and the role of waterbirds in the energy and nutrient flow in food webs is needed. Our study demonstrated that sympatric species can have a trophic plasticity, being highly adaptable to different conditions and resource availability, sharing foraging areas with their conspecifics, indicating important and complex ecological interactions during the breeding season.

Acknowledgements The authors are grateful to colleagues from Waterbirds and Sea Turtles Laboratory, and Institute of Biological Sciences-FURG for fieldwork, data analysis, and laboratory support, as well as contributions to previous drafts of this manuscript, especially Aline B. Silva, Bruno A. Linhares, Cheyenne Soler, Felipe C. Silva, Fernando A. Faria, Juliana V. Gaiotto and Vinícius S. Domingues. We also thank

FURG technicians for assistance in laboratory analysis, Márcio A. Freire for assistance in the identification of fish otoliths, and Leandro R. Ruiz for authorization to study in his property. The authors are also grateful to D. Gianuca and Alexandre M. Garcia for insightful comments on a previous version of this manuscript; and Centro Nacional de Pesquisa e Conservação de Aves Silvestres (CEMAVE) and SISBIO (ICMBio) for bird bands and sampling permits. A. Travessas held a fellowship from the Coordination of Improvement of Higher Education Personnel (CAPES). L. Bugoni is a CNPq research fellow (Proc. No. 310145/2022-8).

Funding The Article Processing Charge (APC) for the publication of this research was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) (ROR identifier: 00x0ma614). This work was funded by the National Council for Scientific and Technological Development-CNPq (Grants PQ No. 310145/2022-8 and Universal 422759/2016-3 to L.B.) and Coordination of Improvement of Higher Education Personnel-CAPES (Finance Code 001).

Data availability Enquiries about data availability should be directed to the authors.

Declarations

Conflict of interest The authors have not disclosed any competing interests.

Ethical and sampling approval This study was approved by ICMBio/SISBIO (sampling permit No. 72947-1) and the Ethics Committee on Animal Use (CEUA-FURG, Certificate No. P028/2020).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abdullahm, M., R. A. Khan, M. Rafay, T. Hussain, T. Ruby, F. Rehman, S. Khalil & S. Akhtar, 2017. Habitat ecology and breeding performance of cattle egret (*Bubulcus ibis*) in Faisalabad, Pakistan. *Pakistan Journal of Zoology* 49:

- 1863–1870. <https://doi.org/10.17582/journal.pjz/2017.49.5.1863.1870>.
- Antón-Tello, M., V. O. Britto, J. A. Gil-Delgado, E. Rico, J. I. Dies, J. S. Monrós & P. Vera, 2021. Unravelling diet composition and niche segregation of colonial waterbirds in a Mediterranean wetland using stable isotopes. *Ibis* 163: 913–927. <https://doi.org/10.1111/ibi.12928>.
- Ashoori, A., H. V. Moradi, H. R. Rezaiee & A. S. Mahiny, 2017a. Dietary segregation of four ardeid species breeding in Anzali International Wetland, northern Iran. *Waterbirds* 40: 377–389. <https://doi.org/10.1675/063.040.0409>.
- Ashoori, A., H. V. Moradi, H. R. Rezaiee & A. S. Mahiny, 2017b. Nest position, breeding success and diet of the black-crowned night heron, *Nycticorax nycticorax* in the Anzali Wetland, northern Iran (Aves: Ardeidae). *Zoology in the Middle East* 63: 283–290. <https://doi.org/10.1080/09397140.2017.1349101>.
- Bachir, A. S., C. Barbraud, S. Doumandji & H. Hafner, 2008. Nest site selection and breeding success in an expanding species, the cattle egret *Bubulcus ibis*. *Ardea* 96: 99–107. <https://doi.org/10.5253/078.096.0111>.
- Barreto, C. T., A. Bianchini, C. Morrissey, F. A. Faria & L. Bugoni, 2025. The influence of colony habitat and egg components on lead and cadmium concentrations of great egrets and roseate spoonbills in southern Brazil. *Ecotoxicology* 34: 577–588. <https://doi.org/10.1007/s10646-025-02856-1>.
- Bearhop, S., C. E. Adams, S. Waldron, R. A. Fuller & H. Macleod, 2004. Determining trophic niche width: a novel approach using stable isotope analysis. *Journal of Animal Ecology* 73: 1007–1012. <https://doi.org/10.1111/j.0021-8790.2004.00861.x>.
- Begon, M., C. R. Townsend & J. L. Harper, 2007. *Ecología - de individuos a ecosistemas*. Porto Alegre: Artmed.
- Bell, W. J., 1990. *Searching behavior: the behavioural ecology of finding resources*. Chapman and Hall Animal Behavior Series. Springer, Dordrecht. <https://doi.org/10.1007/978-94-011-3098-1>.
- Bó, N. A. & C. A. Darrieu, 1993. Aves Ciconiformes; Ardeidae, Ciconiidae. In *Fauna de agua dulce de la República Argentina*, PROFADU (CONICET), Buenos Aires, Argentina, 43: 1B.
- Boecklen, W. J., C. T. Yarnes, B. A. Cook & A. C. James, 2011. On the use of stable isotopes in trophic ecology. *Annual Review of Ecology, Evolution, and Systematics* 42: 411–440. <https://doi.org/10.1146/annurev-ecolsys-102209-144726>.
- Bond, A. L. & K. A. Hobson, 2012. Reporting stable-isotope ratios in ecology: recommended terminology, guidelines and best practices. *Waterbirds* 35: 324–331. <https://doi.org/10.1675/063.035.0213>.
- Britto, V. O. & L. Bugoni, 2015. The contrasting feeding ecology of great egrets and roseate spoonbills in limnetic and estuarine colonies. *Hydrobiologia* 744: 187–210. <https://doi.org/10.1007/s10750-014-2076-1>.
- Brown, S., J. Bizarro, G. Cailliet & D. Ebert, 2012. Breaking with tradition: redefining measures for diet description with a case study of the Aleutian skate *Bathyraja aleutica* (Gilbert 1896). *Environmental Biology of Fishes* 95: 3–20. <https://doi.org/10.1007/s10641-011-9959-z>.
- Brussee, B. E., P. S. Coates, I. A. Dwight & L. G. Young, 2017. Observations of indirect filial cannibalism in response to nest failure of black-crowned night-herons (*Nycticorax nycticorax*). *Wilson Journal of Ornithology* 129: 390–394. <https://doi.org/10.1676/16-013.1>.
- Brzorad, J. N. & A. D. Maccarone, 2013. Activity patterns of snowy egrets (*Egretta thula*) and great egrets (*Ardea alba*): a seasonal comparison. *Waterbirds* 36: 11–19. <https://doi.org/10.1675/063.036.0104>.
- Bugoni, L. & C. M. Vooren, 2004. Feeding ecology of the common tern *Sterna hirundo* in a wintering area in southern Brazil. *Ibis* 146: 438–453. <https://doi.org/10.1111/j.1474-919X.2004.00277.x>.
- Cardarelli, E., M. Fasola, A. Martinoli & D. Pellitteri-Rosa, 2017. Long-term changes in food intake by grey herons (*Ardea cinerea*), black-crowned night-herons (*Nycticorax nycticorax*) and little egrets (*Egretta garzetta*) foraging in rice fields in Italy. *Waterbirds* 40: 344–352. <https://doi.org/10.1675/063.040.0406>.
- Caseiro-Silva, F., F. A. Faria, C. T. Barreto, C. N. Fernandez & L. Bugoni, 2023. Colonial waterbirds provide persistent subsidies to swamp forests along an estuarine island food chain. *Oecologia* 202: 113–127. <https://doi.org/10.1007/s00442-023-05377-y>.
- Castro, D. M. P., D. R. Carvalho, P. D. S. Pompeu, M. Z. Moreira, G. B. Nardoto & M. Callisto, 2016. Land use influences niche size and the assimilation of resources by benthic macroinvertebrates in tropical headwater streams. *PLoS One* 11: e0150527. <https://doi.org/10.1371/journal.pone.0150527>.
- Cherel, Y., K. A. Hobson & S. Hassani, 2005. Isotopic discrimination between food and blood and feathers of captive penguins: implications for dietary studies in the wild. *Physiological and Biochemical Zoology* 78: 106–115. <https://doi.org/10.1086/425202>.
- Cloyed, C. S., K. P. DaCosta, M. R. Hodanbosi & R. H. Carmichael, 2020. The effects of lipid extraction on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values and use of lipid-correction models across tissues, taxa and trophic groups. *Methods in Ecology and Evolution* 11: 751–762. <https://doi.org/10.1111/2041-210X.13386>.
- Comte, L., J. Cucherousset, S. Boulêtreau & J. D. Olden, 2016. Resource partitioning and functional density of worldwide freshwater fish communities. *Ecosphere* 7: e01356. <https://doi.org/10.1002/ecs2.1356>.
- Conversani, V. R. M., M. R. Brenha-Nunes, C. Santificetur, M. B. Giaretta, C. C. Siliprandi & C. L. D. B. Rossi-Wongtschowski, 2017. Atlas of marine bony fish otoliths (sagittae) of southeastern-southern Brazil Part VII: Atheriniformes, Beloniformes, Beryciformes, Zeiformes, Syngnathiformes, Scorpaeniformes and Tetraodontiformes. *Brazilian Journal of Oceanography* 65: 400–447. <https://doi.org/10.1590/S1679-87592017134306503>.
- Craig, E. C., S. B. Elbin, J. P. Sparks & P. D. Curtis, 2015. Identifying important foraging habitat for colonial waterbirds in an urban estuary: a stable isotope approach. *Waterbirds* 38: 330–338. <https://doi.org/10.1675/063.038.0410>.
- Davis-Jr., W. E., 1993. Black-crowned night-heron (*Nycticorax nycticorax*). In: Poole, A. & F. Gill. (eds) *The birds*

- of North America, No. 74. The Academy of Natural Sciences, Philadelphia/The American Ornithologists' Union/Washington, DC.
- Day-Jr, J. W., A. Yáñez-Arancibia, W. M. Kemp & B. C. Crump, 2013. Introduction to estuarine ecology. In: Day-Jr, J. W., B. C. Crump, W. M. Kemp & A. Yáñez-Arancibia (eds). Estuarine ecology. Hoboken: Wiley-Blackwell, p. 1–18. <https://doi.org/10.1002/9781118412787.ch1>.
- Dias, R. A., G. N. Maurício & L. Bugoni, 2017. Birds of the Patos Lagoon Estuary and adjacent coastal waters, southern Brazil: species assemblages and conservation implications. *Marine Biology Research* 13: 108–120. <https://doi.org/10.1080/17451000.2016.1209525>.
- Dias, R. M., E. A. L. Kashiwaqui, J. C. B. Silva, H. Ortêncio-Filho, L. C. Gomes, G. T. R. Souza, R. M. Tófoli, M. H. Machado & A. A. Agostinho, 2025. Feeding ecology of the sympatric waterbirds in Neotropical floodplain. *Hydrobiologia* 852: 751–763. <https://doi.org/10.1007/s10750-024-05674-4>.
- Ducommun, M. P., M. A. Quiroga, A. B. Beltzer & J. A. Schnack, 2008. Diet of cattle egrets in the flood valley of the Paraná River, north Argentina. *Avian Biology Research* 1: 145–151. <https://doi.org/10.3184/175815508X4024>.
- Elliott, K. H., J. D. Roth & K. Crook, 2017. Lipid extraction techniques for stable isotope analysis and ecological assays. In: Bhattacharya, S. K. (ed). *Lipidomics, methods and protocols*, p. 9–24. New York: Humana Press. https://doi.org/10.1007/978-1-4939-6996-8_2.
- Faria, F. A., A. Silva-Costa, D. Gianuca & L. Bugoni, 2016. Cooi heron (*Ardea cocoi*) connects estuarine, coastal, limnetic and terrestrial environments: an assessment based on conventional dietary and stable isotope analysis. *Estuaries and Coasts* 39: 1271–1281. <https://doi.org/10.1007/s12237-016-0073-5>.
- Faria, F. A., E. F. Albertoni & L. Bugoni, 2018. Trophic niches and feeding relationships of shorebirds in southern Brazil. *Aquatic Ecology* 52: 281–296. <https://doi.org/10.1007/s10452-018-9663-6>.
- Fischer, L. G., L. E. D. Pereira & J. P. Vieira, 2004. *Peixes estuarinos e costeiros*, 2nd. edn., Rio Grande: Luciano Gomes Fischer.
- Fry, B., 2006. *Stable isotope ecology*, Springer, New York: <https://doi.org/10.1007/0-387-33745-8>.
- Garcia, A. M., J. P. Vieira & K. O. Winemiller, 2001. Dynamics of the shallow-water fish assemblage of the Patos Lagoon estuary (Brazil) during cold and warm ENSO episodes. *Journal of Fish Biology* 59: 1218–1238. <https://doi.org/10.1111/j.1095-8649.2001.tb00187.x>.
- Garcia, A. M., D. J. Hoeninghaus, J. P. Vieira & K. O. Wine-miller, 2007. Isotopic variation of fishes in freshwater and estuarine zones of a large subtropical coastal lagoon. *Estuarine, Coastal and Shelf Science* 73: 399–408. <https://doi.org/10.1016/j.ecss.2007.02.003>.
- Gause, G. F., 1937. Experimental populations of microscopic organisms. *Ecology* 18: 173–179. <https://doi.org/10.2307/1930461a>.
- Gianuca, D., S. R. Klein & C. M. Vooren, 2010. Dieta de três espécies de garças durante a estação reprodutiva no estuário da Lagoa dos Patos. In: *Anal. of the IV Congresso Brasileiro de Oceanografia* pp. 1–3. Rio Grande: AOCEANO.
- Green, A. J. & J. Elmberg, 2014. Ecosystem services provided by waterbirds. *Biological Reviews* 89: 105–122. <https://doi.org/10.1111/brv.12045>.
- Haimovici, M. & L. G. Cardoso, 2016. Long-term changes in the fisheries in the Patos Lagoon estuary and adjacent coastal waters in southern Brazil. *Marine Biology Research* 13: 135–150. <https://doi.org/10.1080/17451000.2016.1228978>.
- Hamada, N., J. H. Thorp & D. C. Rogers (eds), 2018. *Thorp and Covich's freshwater invertebrates: Volume 3: Keys to Neotropical Hexapoda*. Academic Press.
- Hancock, J. & J. A. Kushlan, 2010. *The herons handbook*, A&C Black, London:
- Herzka, S. Z., E. Mellink, D. M. Talley, G. R. Huxel & P. K. Dayton, 2013. Stable isotope ratios of egg albumen of three waterbird species nesting in the Colorado River Delta indicate differences in foraging ground and isotopic niche breadth. *Aquatic Conservation: Marine and Freshwater Ecosystems* 23: 546–563. <https://doi.org/10.1002/aqc.2326>.
- Hette-Tronquart, N., 2019. Isotopic niche is not equal to trophic niche. *Ecology Letters* 5: 10–12. <https://doi.org/10.1111/ele.13218>.
- Horn, H. S., 1966. Measurement of “overlap” in comparative ecological studies. *American Naturalist* 100: 419–424. <https://doi.org/10.1086/282436>.
- Hothem, R. L., B. E. Brussee, W. E. Davis-Jr., A. Martínez-Vilalta, A. Motis & G. M. Kirwan, 2020. Black-crowned night heron (*Nycticorax nycticorax*), version 1.0. In: Billerman, S. M. (ed). *Birds of the world*. Ithaca: Cornell Lab of Ornithology. <https://doi.org/10.2173/bow.bcnher.01>.
- Hurlbert, S. H., 1978. The measurement of niche overlap and some relatives. *Ecology* 59: 67–77. <https://doi.org/10.2307/1936632>.
- Hutchinson, G. E., 1957. Concluding remarks. *Cold Spring Harbor Symposia on Quantitative Biology* 22: 415–427. <https://doi.org/10.1101/SQB.1957.022.01.039>.
- INMET, 2023. Banco de dados meteorológicos: Dados históricos anuais para município de Rio Grande, Rio Grande do Sul, Instituto Nacional de Meteorologia. <https://portal.inmet.gov.br/dadoshistoricos>. Accessed 11 October 2021.
- Jackson, A. L., R. Inger, A. C. Parnell & S. Bearhop, 2011. Comparing isotopic niche widths among and within communities: SIBER—Stable Isotope Bayesian Ellipses in R. *Journal of Animal Ecology* 80: 595–602. <https://doi.org/10.1111/j.1365-2656.2011.01806.x>.
- Kaufman, K., 1996. *Lives of North American birds*. Houghton Mifflin Company. 2001st ed. Boston.
- Kazantzidis, S. & V. Goutner, 2005. The diet of nestlings of three Ardeidae species (Aves, Ciconiiformes) in the Axios Delta, Greece. *Belgian Journal of Zoology* 135: 165–170.
- Kennish, M. J., 2019. *Ecology of estuaries v. 2: biological aspects*. Boca Raton: CRC Press. <https://doi.org/10.1201/9781351071604>.
- Kioko, J., E. Boyd, E. Schaeffer, S. Tareen & C. Kiffner, 2016. Cattle egret *Bubulcus ibis* interactions with large mammals in the Tarangire-Manyara ecosystem, Northern Tanzania. *Scopus, Journal of East African Ornithology* 36: 15–20.

- Klein, A. H. F. 1997. Regional climate. In: Seeliger, U., C Odebrecht & J. P. Castello (eds). Subtropical convergence environments: the coast and sea in the southwestern Atlantic. Springer, Berlin, Heidelberg, p. 5–7.
- Kushlan, J. A. & J. A. Hancock, 2005. The herons, Oxford University Press, Oxford.
- Layman, C. A., M. S. Araujo, R. Boucek, C. M. Hammer-schlag-Peyer, E. Harrison, R. J. Zachary, P. Matich, A. E. Rosenblatt, J. J. Vaudo, L. A. Yeager, D. M. Post & S. Bearhop, 2012. Applying stable isotopes to examine food-web structure: an overview of analytical tools. *Biological Reviews* 87: 545–562. <https://doi.org/10.1111/j.1469-185X.2011.00208.x>.
- Lemos, V. M., M. Lanari, M. Copertino, E. R. Secchi, P. C. O. V. Abreu, J. H. Muelbert, A. M. Garcia, F. C. Dumont, E. Muxagata, J. P. Vieira, A. Colling & C. Odebrecht, 2022. Patos Lagoon estuary and adjacent marine coastal biodiversity long-term data. *Earth System Science Data* 14: 1015–1041. <https://doi.org/10.5194/essd-14-1015-2022>.
- Maccarone, A. D., R. E. Renken & B. L. Hamilton, 2015. Effect of water level and time of day on black-crowned night-heron (*Nycticorax nycticorax*) foraging behavior at an artificial weir: 2013–2014. *Transactions of the Kansas Academy of Science* 118: 90–96. <https://doi.org/10.1660/062.118.0110>.
- Magurran, A. E., 2019. Medindo a diversidade biológica. Curitiba: Editora UFPR.
- Malabarba, L. R., P. Carvalho-Neto, V. D. A. Bertaco, T. P. Carvalho, J. D. Santos & L. G. S. Artioli, 2013. Guia de identificação dos peixes da bacia do rio Tramandaí. Porto Alegre: Via Sapiens.
- Martínez, C., 2010. Trophic niche breadth and overlap of three egret species in a Neotropical mangrove swamp. *Waterbirds* 33: 285–292. <https://doi.org/10.1675/063.033.0303>.
- Martínez-Vilalta, A. & A. Motis, 1992. Family Ardeidae (herons). In del Hoyo, J., A. Elliot & J. Sargatal (eds), *Handbook of the birds of the world (hoatzin to auks)*, Vol. 1. Lynx Editions, Barcelona: 376–429.
- Mittelbach, G. G. & B. J. McGill, 2019. *Community ecology*, 2nd ed. Oxford University Press, Oxford: <https://doi.org/10.1093/oso/9780198835851.001.0001>.
- Morales-Silva, E. & S. N. Del Lama, 2014. Colonization of Brazil by the cattle egret (*Bubulcus ibis*) revealed by mitochondrial DNA. In: Capdevila-Argüelles, L. & B. Zilletti (eds). *Proceedings of 7th NEOBIOTA conference*, Pontevedra, Spain. *NeoBiota* 21: 49–63. <https://doi.org/10.3897/neobiota.21.4966>.
- Navarro, A. B., J. A. Bogoni, M. Z. Moreira & L. F. Silveira, 2023. Intraguild niche partitioning in granivorous birds from the late past. *Avian Research* 14: 100075. <https://doi.org/10.1016/j.avrs.2023.100075>.
- Newsome, S. D., C. Martínez del Rio, S. Bearhop & D. L. Phillips, 2007. A niche for isotopic ecology. *Frontiers in Ecology and the Environment* 5: 429–436. <https://doi.org/10.1890/060150.1>.
- Nunes, M. F., A. L. Roos, R. C. Barbosa-Filho & L. A. Mes-tre, 2010. The cattle egret (*Bubulcus ibis*) on Fernando de Noronha Archipelago: history and population trends. *Revista Brasileira de Ornitologia* 18: 315–327.
- Park, S.-R., K.-Y. Kim, H. Chung, Y.-S. Choi & H.-C. Sung, 2011. Vertical nest stratification and breeding success in a six mixed-species heronry in Taeseong, Chungbuk, Korea. *Animal Cells and Systems* 15: 85–90. <https://doi.org/10.1080/19768354.2011.555119>.
- Parnell, A. C., D. L. Phillips, S. Bearhop, B. X. Semmens, E. J. Ward, J. W. Moore & R. Inger, 2013. Bayesian stable isotope mixing models. *Environmetrics* 24: 387–399. <https://doi.org/10.1002/env.2221>.
- Parsons, K. C. & T. L. Master, 2020. Snowy egret (*Egretta thula*), version 1.0. In: Poole, A. F. & F. B. Gill (eds). *Birds of the world*. Ithaca: Cornell Lab of Ornithology. <https://doi.org/10.2173/bow.snoegr.01>.
- Pereira, N. & F. D’Incao, 2012. Relationship between rainfall, pink shrimp harvest (*Farfantepenaeus paulensis*) and adult stock, associated with El Niño and La Niña phenomena in Patos Lagoon, southern Brazil. *Journal of Marine Biological Association of the United Kingdom* 92: 1451–1456. <https://doi.org/10.1017/S0025315412000021>.
- Post, D. M., 2002. Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology* 83: 703–718. [https://doi.org/10.1890/0012-9658\(2002\)083\[0703:USITET\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[0703:USITET]2.0.CO;2).
- Post, D. M., C. A. Layman, D. A. Arrington, G. Takimoto, J. Quattrochi & C. G. Montana, 2007. Getting to the fat of the matter: models, methods and assumptions for dealing with lipids in stable isotope analyses. *Oecologia* 152: 179–189. <https://doi.org/10.1007/s00442-006-0630-x>.
- Quintela, F., F. Corrêa, R. M. Pinheiro & D. Loebmann, 2018. Ichthyofauna of Marinheiros Island, Patos Lagoon estuary, southern Brazil. *Biota Neotropica* 18: e20170430. <https://doi.org/10.1590/1676-0611-BN-2017-0430>.
- Quiroga, M. A., E. J. Leon, P. F. Olguin & A. H. Beltzer, 2013. Diet of black-crowned night-herons (*Nycticorax nycticorax*) in a wetland of the Paraná River’s Alluvial Valley. *Ekoloji* 22: 43–50. <https://doi.org/10.5053/ekoloji.2013.886>.
- R Core Team, 2020. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Régale, M. A., D. Siel & M. Ruibe, 2014. Bait fishing black-crowned night heron (*Nycticorax nycticorax*) in Chile. *Ornitología Neotropical* 25: 465–468.
- Riehl, C., 2006. Widespread cannibalism by fledglings in a nesting colony of black-crowned night herons. *Wilson Journal of Ornithology* 118: 101–104. <https://doi.org/10.1676/04-119.1>.
- Ruiz-Guerra, C. & M. Á. Echeverry-Galvis, 2019. Prey consumed by wading birds in mangrove swamps of the Caribbean coast of Colombia. *Journal of Natural History* 53: 1823–1836. <https://doi.org/10.1080/00222933.2019.1667037>.
- Salas-López, A., C. Violle, F. Munoz, F. Menzel & J. Orivel, 2022. Effects of habitat and competition on niche partitioning and community structures in Neotropical ants. *Frontiers in Ecology and Evolution* 10: 863080. <https://doi.org/10.3389/fevo.2022.863080>.
- Sánchez-García, I., 2012. Dieta herpetófaga en una garcilla bueyera (*Bubulcus ibis*). *Boletín de la Asociación Herpetológica Española* 23: 10–12.
- Schoener, T. W., 1974. Resource partitioning in ecological communities. *Science* 185: 27–29. <https://doi.org/10.1126/science.185.4145.2>.

- Silva, E. B., M. F. Nóbrega, A. M. Grimm, M. S. Copertino, J. P. Vieira & A. M. Garcia, 2021. Long-term trends in the abundance of an estuarine fish and relationships with *El Niño* climatic impacts and seagrass meadows reduction. *Estuarine, Coastal and Shelf Science* 261: 107565. <https://doi.org/10.1016/j.ecss.2021.107565>.
- Silva-Costa, A. & L. Bugoni, 2013. Feeding ecology of kelp gulls (*Larus dominicanus*) in marine and limnetic environments. *Aquatic Ecology* 47: 211–224. <https://doi.org/10.1007/s10452-013-9436-1>.
- Smith, J. P., 1997. Nesting season food habits of 4 species of herons and egrets at Lake Okeechobee, Florida. *Colonial Waterbirds* 20: 198–220. <https://doi.org/10.2307/1521686>.
- Sovrano, L. V., S. A. Regner, R. E. Lorenzón, G. N. Ceppi, A. Rocha & A. H. Beltzer, 2022. Canibalismo en *Nycticorax nycticorax* y depredación interespecífica de juveniles de *N. nycticorax* sobre pichones *Ardea alba* (Ardeidae) en una colonia mixta en Argentina. *Oecologia Australis* 26: 620–629. <https://doi.org/10.4257/oeco.2022.2604.09>.
- Staddon, P. L., 2004. Carbon isotopes in functional soil ecology. *Trends in Ecology & Evolution* 19: 148–154. <https://doi.org/10.1016/j.tree.2003.12.003>.
- Steinmetz, J., S. L. Kohler & D. A. Soluk, 2003. Birds are overlooked top predators in aquatic food webs. *Ecology* 84: 1324–1328. [https://doi.org/10.1890/0012-9658\(2003\)084\[1324:BAOTPI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2003)084[1324:BAOTPI]2.0.CO;2).
- Teixeira-Amaral, P., W. J. A. Amaral & D. O. Ortiz, 2017. The mesozooplankton of the Patos Lagoon Estuary, Brazil: trends in community structure and secondary production. *Marine Biology Research* 13: 48–61. <https://doi.org/10.1080/17451000.2016.1248850>.
- Telfair-II, R. C., 2023. Western cattle egret (*Bubulcus ibis*), version 1.0. In: Rodewald, P. G., B. K. Keeney, S. M. Billerman & M. A. Bridwell (eds). *Birds of the world*. Ithaca: Cornell Lab of Ornithology. <https://doi.org/10.2173/bow.cater1.01>.
- Vega-Sánchez, V., C. I. Lomelí-Chávez, J. A. Montaña-Reyes, N. E. Reyes-Rodríguez, F. R. Gómez-de Anda, N. L. Calderón-Apodaca & A. P. Zepeda-Velázquez, 2022. Elements that make up the diet of the cattle egret (*Bubulcus ibis*) in Hidalgo, Mexico. *Huitzil* 23: e-632. <https://doi.org/10.28947/hrmo.2022.23.1.636>.
- Wagner, D. L., E. M. Grames, M. L. Forister, M. R. Berenbaum & D. Stopak, 2021. Insect decline in the Anthropocene: death by a thousand cuts. *Proceedings of the National Academy of Sciences of the United States of America* 118: e2023989118. <https://doi.org/10.1073/pnas.2023989118>.
- Weller, M. W., 1999. *Wetland birds: habitat resources and conservation implications*, Cambridge University Press, Cambridge.
- Williams, C. T., C. L. Buck, J. Sears & A. S. Kitaysky, 2007. Effects of nutritional restriction on nitrogen and carbon stable isotopes in growing seabirds. *Oecologia* 153: 11–18. <https://doi.org/10.1007/s00442-007-0717-z>.
- Williamson, L., M. Hudson, M. O'Connell, N. Davidson, R. Young, T. Amano & T. Székely, 2013. Areas of high diversity for the world's inland-breeding waterbirds. *Biodiversity and Conservation* 22: 1501–1512. <https://doi.org/10.1007/s10531-013-0488-2>.
- Ye, Y., C. Hu, Y. Jiang, G. W. Davison & C. Ding, 2021. Three-dimensional niche partitioning between two colonially nesting ardeid species in central China. *Avian Research* 12: 33. <https://doi.org/10.1186/s40657-021-00264-7>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.